

Color K

POTASSIUM FERTILIZER.



COMPOSITION	% w/v
Potassium (K ₂ O)	32,0
EDTA	2,0
pH	12 - 13
Density	1,25

Improves the flavour of fruit. Raises the sugar content and increases the consistency of the fruit. Especially indicated during swelling and ripening.

Applied to olive trees, it improves the fat content. The presence of EDTA increases efficiency by improving the availability of potassium in the plant when it most needs it. **RECOMMENDED FOR ALL TYPES OF CROPS.**

CROPS		DOSAGE
Cotton:	2-4 treatments during the crop's life cycle.	<p>FOLIAR DOSAGE: 130-350 cc / 100 Lts</p> <p>FERTIRRIGATION DOSAGE: 40-60 Lts / Ha every 15 days.</p>
Citrus Fruits:	Apply when the fruit is setting, swelling and before harvesting.	
Strawberries:	1-3 treatments during flowering, fruit formation and formation of the tubers.	
Fruit Trees:	Apply when the fruit is setting, swelling and before harvesting.	
Tropical Fruits:	2-4 treatments during the crop's life cycle.	
Horticultural Crops:	2-6 applications throughout the crops vegetative cycle.	
Olive Trees:	Apply when the fruit is setting, swelling and before harvesting.	
Sugar Beet:	From 2 months before harvesting and onwards.	
Grapes:	Apply when the fruit is swelling, ripening and gaining colour.	

PRESENTATION

