

	SOIL DOSAGE	Lts/ha
STRAWBERRIES	Every 10 days after transplanting	4
FRUIT TREES	From budding until the swelling of the fruit	6
BANANA PLANTS	Every 15 days between March and June	6
OLIVE TREES	Throughout the whole cycle	18
TABLE GRAPES	From budding until the end of the cycle	5
DRY FRUITS	From budding until the swelling of the fruit	5
CITRUS FRUIT	From flowering until the swelling of the fruit	-
COTTON	10 days after shooting until 20 days after the flowering	12
ORNAMENTAL PLANTS	Every 15 days after transplanting	6
	· · · · •	4

	FOLIAR DOSAGE	cc/100L
HORTICULTURAL CROPS STRAWBERRIES TUBERS FRUITTREES BANANA PLANTS OLIVETREES TABLE GRAPES WINE GRAPES DRY FRUITS CITRUS FRUITS BEET COTTON ALFALFA ORNAMENTAL PLANTS LAWN	Every 10 days after transplanting Throughout the whole cycle Every 15 days From budding until the swelling of the fruit Every 15 days Throughout the whole cycle From budding until the end of the cycle From budding until the end of the cycle From budding until the swelling of the fruit From flowering until the swelling of the fruit 2 applications every 15 days 10 days after sprouting until 20 days after the first flower. After every mowing Every 15 days after transplanting After sowing/Growth phase	200 200 250 250 200-300 250 2L/Ha 200-300 200-300 200-300 2,5L/Ha 300 2,5L/Ha 250 3-5L/Ha/30.cc/mi

## AMINOACIDS

COMPOSITION % w/w Free Amino Acids Total Nitrogen(N) (N organic) Organic Carbon Total Organic Material ISI (Disease-Resistance Activator) C/N Ratio PH 5,6 Density 1,25	25,00 6,4 27,5 47,00 3,00 4,30	
Inorganic Nitrogen (ammonium chloride) Absent Vitamins 0,96		

STYM 25 is a natural bioactivator based on Amino acids obtained through enzymatic processes, making STYM25 more efficient than chemicalprocess based products. It is recommended for all crops and all times, especially when the plants need more nutrients such as in pre-floration, setting, the swelling of the fruit, vegetative growth, for saline or climatic conditions etc. Agrícola Aspe has developed a group of molecules that we call I.S.I. capable of acting as DISEASE RESISTANCE ACTIVATORS.

