

NEW
Imported from Spain

AF Gibberellic SL

PLANT GROWTH REGULATOR

AF Gibberellic is a plant growth regulator characterized by their physiological and morphological effects. Acts at very low concentrations; is translocated inside the the plant and usually affects only the aerial parts.

AF Gibberellic reinforces apical dominance stimulating flowering, fruiting set, breaking the dormancy of seeds and vegetative organs and removing stress from some virus.

CAUTIONS FOR USE

Ensure that the wind does not drag the spraying to other neighboring crops. Apply the product shortly after mixing with water to prevent decomposition.

In the event of a precipitation at 8 hours after

USAGE INSTRUCTIONS:

Treat with high water volume (600-1500 l / ha).

Add wetting

COMPOSITION


Gibberellic acid (GA3)
1,6% (w/v)
Soluble liquid (SL)



**GROWTH
BLOOMING
AND FRUIT SET**

CROPS/TREATMENT	DOSES
Clementine: To improve the consistency of the peel (when the green treat is gone).	30-50 cc/hl
Lemon and Clementine boneless: To induce fruit set and fruit setting. Treat at petal drop and repeat the treatment at 3-4 weeks.	40 cc/hl
Tangerine: To prevent fruit drop (treat at petal drop, repeating at 3-4 weeks) and improve the consistency of the peel (when the green treat is gone).	40 cc/hl
Pear (cv. Blanquilla): To reduce fruit drop during filling and prevent frost damage. Treat with 30-60% open flower or 48 hours to avoid the effect of frost.	60-70 cc/hl
Vine (cv. Macabeo): To induce the elongation of the cluster and peduncles of the fruit. Apply before falling flowerhoods.	30-50 cc/hl
Strawberry: To promote fruit set and fruit growth. Treat a little before the start of flowering. Can be reduced the dose in plantations over a year.	60 cc/hl

Packaging



NON TOXIC

Aspe