

NEW
Imported from Spain

AZOL

NITROGEN FERTILIZER OF CONTROLLED RELEASE

CHARACTERISTICS

AZOL is a nitrogen fertilizer of high efficiency and persistence. The **N-urea** meet immediate needs and the **N-formaldehyde** ensures progressive availability. Safe and environmentally friendly.

GENERAL ADVANTAGES

Progressive release.
Compatible with pesticides.
Foliar application.
Reduce, drift and improves the absorption.
Keep vegetative/productive balance in crops where (N) has direct influence on production.
Risk of washing, maximum use of nitrogen, environmentally friendly.

CEREAL

Specific weight increases
Increases protein content
Decreases risk of lodging

BENEFITS CROPS

MAIZE

In the early stages homogenize the crops.

RICE

Increases the specific weight . Largest filled of the Spike.

OLIVE

Homogenized fruit ripening.
Greater fruit size.
Net increase production.

COMPOSITION

%w/v

Total Nitrogen (N)	28%
• N (Urea-formaldehyde)	85%
• N (Ureico)	15%

Biuret, Max	less than	0,7%
pH(water 10%, 20°C)		9-11
Density (20°)		1,23-1,25kg/L



CROPS	DAYS	STATE OR USE MODE	DOSAGE
Cotton	n.p.	In the phenological stages of 4 sheets, after flowering and during capsule formation	5-8 l/ha - 6-10 kg/ha
Rice	n.p.	The emergence of the flag leaf	10-15l/ha - 12-19 kg/ha
Cereals	n.p.	1st treatment between full tillering to the first knot. 2nd treatment between flag leaf and flowering	10-25 l/ha - 12-31 kg/ha
Lawn	n.p.	Applications every 50-60 days	15-24 l/ha - 19-30 kg/ha
Citrus	n.p.	After fruit set and floral differentiation	5-10 l/ha - 6-12 kg/ha
Citrus	n.p.	End of winter, from vegetative recovery	64-96 l/ha - 80-120 kg/ha fertirrigation
large strawberries	n.p.	Beginning of flowering and repeat every 15 days	5-10 l/ha - 6-12 kg/ha
Stone fruit	n.p.	Fruit before flowering and repeat every 30 days	5-10 l/ha - 6-12 kg/ha
Stone fruit	n.p.	End of the winter to fruit set	64-96 l/ha - 80-120 kg/ha fertirrigation
Pome fruit	n.p.	Before flowering and repeat every 30 days	5-10 l/ha - 6-12 kg/ha
Pome fruit	n.p.	End of the winter to fruit set	64-96 l/ha - 80-120 kg/ha fertirrigation
Leafy vegetables	n.p.	The formation of the bud and repeat every 14 days	5-8 l/ha - 6-10
Horticultural	n.p.	Before flowering and applications every 10-14 days	5-10 l/ha 6-12 kg/ha
Horticultural	n.p.	At the beginning of the growing season	48-80 l/ha 60-100 kg/ha fertirrigation
Corn	n.p.	Early post-emergence, together with herbicides	10-15 l/ha - 12-19 kg/ha
Corn	n.p.	Acaricide treatment for spider mites	10-15 l/ha - 12-19 kg/ha
Olive	n.p.	Since before flowering and repeat every 30 days	5-10 l/ha - 6-12 kg/ha
Olive	n.p.	End of winter, from vegetative recovery	64-120l/ha - 80-150 kg/ha fertirrigation
Soy	n.p.	State R2-R3	4-8 l/ha - 5-10 kg/ha
Vine	n.p.	Post-flowering	5-10 l/ha - 6-12 kg/ha
Vine	n.p.	End of winter, from vegetative recovery	48-80 l/ha - 60-100 kg/ha fertirrigation

Packing



Aspe